

## Work-related stress and depressive disorders

Christopher Tennant\*

*Department of Psychological Medicine, University of Sydney and Royal North Shore Hospital, Block 4, level 5, St. Leonards, Sydney, NSW 2065, Australia*

*Institute of Stress Medicine, University of Sydney and Royal North Shore Hospital, Block 4, level 5, St. Leonards, Sydney, NSW 2065, Australia*

Received 26 May 1999; accepted 13 March 2001

### Abstract

The 1980s and 1990s has seen a considerable change in the workforce structure in industrialised economies. Employees are commonly faced with greater demands and less job security, both of which are likely to be stressful, thus psychological disorders especially depression may increasingly be caused by work-related stressors. An issue of this journal in 1997 (Vol. 43, No. 1) was indeed devoted to stress in the workplace and since then, these workplace changes have progressed and a review seems timely. Because interpreting results of cross-sectional studies is limited by a potential reciprocal relation between work stressors and depression (since “effort after meaning” can influence how “distressed”

individuals report stressors at work), this review largely focuses on prospective or predictive studies to minimise this bias. Not surprisingly, the findings from occupational stress research is consistent with the more general life event stress literature showing that specific acute work-related stressful experiences contribute to ‘depression’ and, more importantly perhaps, that enduring ‘structural’ occupational factors, which may differ according to occupation, can also contribute to psychological disorders. There are significant implications for employees, their families, employers and indeed the wider community. © 2001 Elsevier Science Inc. All rights reserved.

**Keywords:** Work; Stress; Depression; Burnout

### Introduction

Work and family are the two domains from which most adults derive satisfaction in life; equally they are the common sources of stressful experiences. The working environment continues to change with globalisation of the world economy and economic rationalisation driving job restructuring, greater part-time and contract work, and greater workload demands that commonly occur in a context of higher job insecurity. There is thus not an unreasonable perception in the community that work is increasingly the source of much of our stress and distress.

The implications of work-related stress include the effects on worker satisfaction and productivity, their mental and physical health, absenteeism and its economic cost, the wider impact on family function and finally, the

potential for employer liability. While depression is the most likely adverse psychological outcome, the range of other possible “psychological” problems include “burnout,” alcohol abuse, unexplained physical symptoms, “absenteeism,” chronic fatigue and accidents, sick building syndrome and repetitive strain injury [1]. This review focuses largely on prospective or “predictive” studies both of “depression” and “burnout,” and it is based on literature searches using Medline, Psych INFO and EMBASE from 1966 to 2000 (keywords: occupational stress, work, occupations, workload, occupational health, depression, anxiety). The review assesses the stressor findings by different occupational groups because they may embrace different qualities of stress which furthermore may be a major factor contributing to psychological morbidity in these groups.

### Morbidity in different occupational groups

In the studies reviewed it is usual for *depression* to be diagnosed by a range of questionnaires including the

\* Department of Psychological Medicine, University of Sydney and Royal North Shore Hospital, Block 4, level 5, St. Leonards, Sydney, NSW 2065, Australia. Tel.: +61-2-9926-7746, +61-2-9926-7147; fax: +61-2-9926-7730.

E-mail address: [tennant@med.usyd.edu.au](mailto:tennant@med.usyd.edu.au) (C. Tennant).

Beck Depression Inventory, Zung scale, the General Health Questionnaire (GHQ), the Centre for Epidemiological Studies Depression Scale (CESD) or similar clinically validated questionnaires. *Burnout* is also a construct often used and embraces three clusters of symptoms; “emotional exhaustion,” “depersonalisation” (negative, insensitive attributes to clients) and a sense of reduced “personal accomplishment” [2]. While burnout by definition is only assessed in an occupational setting, it may nonetheless be a precursor or correlate of depression [3–5].

Three broad but distinct occupational groups are readily identifiable in the literature. They are *blue-collar/manual workers*, *white-collar/civil servants* and the “*helping*” professions (teachers and healthcare workers). The Karasek “demand–control” model of work-related stress [6] is commonly employed, especially in the first two groups. In this model, the composite construct of “job strain” is defined as a combination of the constructs of (high) *job demands* and (low) *decision making latitude* (or control over workload), both of which are “objectively” derived from descriptions of the nature of work. All three of the above constructs have been assessed in various studies.

#### *Blue-collar workers*

In factory and blue-collar workers studied over 3 years, in Japan ( $N=468$ ) [7,8], “lack of control over work,” “unsuitable jobs” and “poor work place relations” predicted depression (Zung scale) [7]. Furthermore, the latter two work variables also predicted depressive disorder diagnosed clinically by a psychiatrist in a nested cohort of this sample; these relationships were significant even after controlling covariates and initial depressive symptoms [8]. In a combined study of 900 blue- and white-collar industrial workers studied over 5 and 10 years [9], stressors at work predicted mental stress symptoms, poor perceived health and absenteeism; mental stress symptoms in turn predicted smoking, drinking, and absenteeism.

In the blue-collar environment, *noise* can be a particular work stressor. In 2368 blue-collar workers [10], noise exposure in women predicted irritability, somatic complaints, anxiety and depression; in men noise predicted only “post work irritability.” These relationships however were found only in “vulnerable” subjects being those reporting “noise annoyance” (the emotional reaction to noise at exposure). In both sexes noise however contributed to increased work accidents and sickness absence [10]. Similarly, in another setting [11], while traffic noise had no direct effect on psychological morbidity, it was related to “noise annoyance” which was associated with “noise sensitivity” (the physiological reaction to noise). The complex relationships of “noise sensitivity,” “noise annoyance,” “negative affectivity” and depression were

explored in two other longitudinal studies [12]. Noise sensitivity was found to be moderately stable over time, and predicted depression, but in part this relationship was reciprocal.

#### *White-collar workers*

In white-collar workers, there have been several large prospective studies. In a 4-year Finnish study ( $N=15530$ ) [13], records linkage was used to access psychiatric morbidity data (suicides, hospitalisation, prescription of psychotropic drugs). Interpersonal conflict at work predicted “physician-diagnosed” psychiatric morbidity (relative risk=2.18) even after controlling social class, prior health and prior “mental instability/stress” (i.e., neuroticism, hostility, life stress, low self-assurance) [13]. In a 5-year prospective study of 5000 employees drawn from a Danish population register [14], changes in repetitive work, job insecurity, high demands and low support all predicted a single “self-rated” health item.

In the impressive Whitehall II study, (6895 male and 3413 female civil servants) [15–21] psychological morbidity (GHQ score) was predicted both by poor work social supports, and by high job demands [15]. Poor social support and poor “skill discretion” also predicted “psychiatric absenteeism” [15]. These results confirmed earlier cross-sectional findings [16]. Both negative work relationships and the imbalance between work effort required and occupational reward also predicted poor physical, psychological and social functioning (SF36), even after controlling confounding variables. [17]. Furthermore, much of the social class difference that had been found in both “depression” (GHQ) and “well being” (affect balance scale) was explained by the psychosocial (Karasek) work environment [18]. Finally, in a quasi-experimental component of the study, “job insecurity” (the likelihood of redundancy), predicted physical symptoms, body mass index, excessive sleep [19] and, psychological morbidity [20]. Job insecurity did not however predict any change in actual health behaviours [21].

In managerial employees ( $N=442$ ) [22], two of the four work-related stressors that were assessed (work load and role ambiguity) predicted depressive symptoms. This association furthermore was potentiated by “negative affectivity,” but only in females [22]. In a “community cohort” of 1415 twenty-six-year-old men in mixed occupations [23], “nervous strain” (a self-report single item) was predicted by occupational status but not by social class of origin. After accounting for occupational status, and neuroticism, five work tasks made a significant contribution to nervous strain. These were supervision, teaching, driving, skilled machine operation and people contact. Finally, in a community-based study of 795 employed adults [24], work pressure, lack of autonomy and role ambiguity predicted subsequent depression (CESD) but job involvement did not.

*“Caring” professions**Teachers*

In “new” or trainee teachers [25], work demands and poor support predicted depression, (GHQ) over 2 months in one study, while in another [26,27], three self-reported work stress indices predicted postemployment depression (CESD scale) and job dissatisfaction, even when initial depression and personality were controlled. In more experienced teachers [28–30], burnout was predicted by both “work setting” and specific “work stress.” The effect of these variables was however not direct, but operated by influencing both the work setting and work stress at the subsequent assessment [29]. Specific work variables predicting burnout were poor social support outside the organisation, personal characteristics [29] and, “red tape” and “disruptive students” [30]. Similarly, teaching “stressful students” early in the school year predicted poor mental health in the latter part of the year in another sample [31].

A more detailed study of the relation of 12 work stress variables to the three “burnout” subscales (in another teacher sample) [32], found “role conflict” alone predicted “emotional exhaustion,” only two “support variables” predicted “poor accomplishment” while there were no predictors of “depersonalisation.” Finally, in school psychologists [33] all three burnout subscales were predicted by personality (neuroticism), but only “emotional exhaustion” and poor “accomplishment” were predicted by self-reported “work stressors.” The relation between work stressors and burnout was furthermore found to be reciprocal, and when initial burnout was controlled, work stressors did not predict subsequent burnout [33].

*Doctors*

The nature of the speciality and the duration of professional experience influences risk of morbidity in doctors. For instance, in junior doctors [34] low autonomy predicted psychological morbidity while work demands were most predictive in older doctors. In relation to medical specialities for instance, work-related stress and specifically, “low task–role clarity” predicted later depression in emergency medicine residents [35], while in GPs [36], routine work administration, job demands, interference with family and interruptions with work, predicted their negative mental well-being. In anaesthetists [37], communication in the hospital and perceived of control over work were significant in determining job satisfaction and mental well-being. In middle-aged medical graduates generally [38], the effects of psychological job demands, patient demands, physician resources and work control were studied; high job demands were associated with both “work dissatisfaction” and psychological disorder (GHQ) in univariate analyses. In regression analyses, lack of control over work was independently associated with both dissatisfaction and psychological disorder [38].

*Nurses and other health workers*

Healthcare workers generally have psychological morbidity rates higher than the general population; in a large NHS sample in the UK the relative risk of disorder was 1.5 and was most marked in direct care staff and women in particular [39].

In nurses [40], “burnout-related absenteeism,” was predicted by “ambiguity about authority” and perceived lack of social support at work; this study was however limited by a low 43% response rate. In another sample of nurses [41] psychological well-being was predicted by the congruence of work status (full- or part-time) with the desire for that degree of employment.

In youth workers [42], perceptions of job insecurity, of a “controlling work setting” and low social support, predicted burnout; eight other indices of work stress were however not significant predictors. In long-term care staff [43], work load and relationship stress, both predicted depression and physical symptoms over 8 months. Background variables and “variables assessed at time one,” predicted much of the variance both in depression (42%) and in physical symptoms (38%), while problems with supervisor relationships predicted only 13% of the variance in depression and physical symptoms, respectively. Paradoxically, patient care stressors were predictors of better psychological and physical health status. The structure of the work environment (cohesion, autonomy, clarity) however was not predictive [43].

In social workers [44], change in burnout over time was independently predicted by job stress, poor supervisor support, low satisfaction with clients and poor self-esteem; together however they accounted for only 28% of the variance [44]. In child protection workers [45], work environment and work hassles predicted all three subscales of burnout over 18 months, the relationship remaining significant with initial burnout controlled. Finally, in a well-designed albeit, cross-sectional study of mental healthcare workers [46] involving a principal components analysis of work stress, some 28% of variance in “depression” (GHQ score) and 42% in burnout were explained by work stress factors; in particular, poor staffing resources, work overload and carer dissatisfaction were related to depression.

Thus while different occupational groups may differ in the nature of their work environments, the “social environment” (usually conflict in relationships or poor social support), seems to predict depressed mood or burnout in most occupational groups. Also reasonably consistent as a predictor are “client-related” stressors in the caring professions. The structure of individual work environments seems more difficult to measure and less consistently predicts depression, although lack of control over work is a relatively consistent predictor. Overall, the proportion of variance in depression/burnout explained by stress factors in different studies varies considerably from approximately 10% to 50%, due in all probability to the

differences in groups studied, the measures used and the varying time frames of assessments.

### “Moderating variables”

Covariates are often presumed to influence the “work stress and depression” relationship; these include specific acute stressful events, “hours worked,” job involvement, job controllability, personality variables and social supports. Whether these variables are independent predictors of depression or moderate the relation of work stress to depression is not always clear.

#### *Specific stressful events in the workplace*

Occupations of different types may be associated with specific types of stressful life events that contribute to psychological disorder. These include negative work relationships or interpersonal conflicts [13,17,43], general abuse and sexual harassment [47]), bullying (especially by supervisors) ([48] and other adverse events including trauma exposure in nurses [49]), air traffic incidents in pilots [50] or stressful and disruptive students for teachers [30,31]. These findings are consistent with the general life events literature. [51]

#### *Hours worked*

The effect of hours worked has been reviewed by Spurgeon et al. [52] who carefully distinguished this variable both from shift work and work overload; 10 of 11 studies revealed an adverse effect of long hours of work on mental health. This relationship, in turn, was further moderated by other variables including employee choice [53], personality [54], female sex [55] and occupational class [56]. Long hours, however, may be employer or employee determined. In relation to employee factors and the impact on psychological health, that of the three components of “workaholism” studied [57] (job involvement, “personal drivenness” and lack of enjoyment), only the latter two variables correlated with poor health indices in MBA graduates.

#### *Job involvement*

In a study of 795 employed adults in a community sample [24], there was only limited support for job involvement having any stress moderating influence on the relationships between role ambiguity and both physical health and alcohol use. There was no evidence for a moderating effect on depression (CESD) [24]. In two earlier studies, one in nurses [40] found some moderating effect while another in supermarket managers [58] did not. Finally in the above study of “workaholism” [57], “job

involvement” did not correlate with measures of psychological distress/well-being.

#### *Job controllability*

In a review of the evidence, Glass and McKnight [2] concluded the “uncontrollability” contributed to negative psychological states largely when there was a “discrepancy between career aspirations and occupational achievement” (i.e., there was career dissatisfaction). This conclusion may, however, be somewhat limited being based largely on findings from cross-sectional studies.

#### *Personality*

Personality factors studied in particular include “negative affectivity” or neuroticism. Some studies show that personality has an *independent effect* on depression and does not simply moderate the relationship of stressors with psychological morbidity. For instance, in a large community sample [23] both neuroticism and specific work stressors separately contributed to a “nervous strain variable”; there was no evidence of an interaction between these variables nor was there evidence that stressful jobs were held particularly by anxiety-prone subjects [23]. In helping professionals [42], the attributional style of “accepting personal responsibility for client outcomes” was also an independent predictor of burnout. Finally, in school psychologists [33] personality variables (including neuroticism) remained significant predictors of burnout (accounting for up to 18% of the variance in some instances) even after controlling for demographic variables and self-reported work stressors.

A “moderating” effect for personality has however been reported in two studies. “Negative affectivity” was found to potentiate the association between work stressors and depression in female (but not male) managerial employees [22] and in teachers [25]. Finally, negative affectivity/neuroticism may also *confound* the relation between stressors and depression because this “variable” is firstly an independent risk factor for emotional symptoms (across most situations and even in the absence of stressors) [59] and secondly because it may contribute to various occupational stress factors including self-reports of stressors [60,61], interpersonal conflicts [62] and work dissatisfaction [63]. Prospective studies best address this latter problem.

#### *Social supports*

Social support is another covariate that may be either a potential moderator or independent risk variable. Poor social support was an *independent predictor* of distress (GHQ) in medical graduates [33], of absenteeism in nurses [40], of psychological symptoms in a community sample of working women [64], and of psychological morbidity (burnout and GHQ score) in Japanese local government officials [65]. In the Whitehall II study [66], social supports at work (even

after adjusting for initial GHQ score) predicted both psychological morbidity (GHQ) and more brief periods of psychiatric absenteeism; this association was further potentiated by poor supports outside work [66]. Poor relationships outside of work and significant marital problems also predicted long periods of psychiatric absenteeism in this sample [67]. Findings overall did not support the stress buffering effect of social supports [66], indeed, paradoxically, good social supports outside of work actually predicted absenteeism (due to both physical and psychiatric problems) [68].

A *buffering effect* has however also been reported. A small prospective study [69] found a general buffering effect on psychological symptoms while two other studies report more restricted impact. For instance, in one study [70] a causal effect was reported in some workers, being only those who had specific and multiple work stressors, while in a community sample [71] only the support of a supervisor reduced the risk of depression over 1 year while support from a colleague did not. As in the literature generally, there remains conflict concerning social support as an independent risk factor for morbidity or as simply a moderator of the relation between stressors and psychological morbidity; the evidence more strongly supports the former.

#### Combined stressors

The effect of stressors of different ‘origins’ (those at work and those outside it) has also been assessed. Both work stress and domestic stress predicted depression both in married professionals (DSM IIIR) even after controlling potential confounders [72] and in female hospital workers [73]. In working women [74], while work overload predicted psychological disorder (GHQ), surprisingly perhaps their family role stressors did not: high occupational status furthermore diminished the impact of work overload on psychological disorder [74].

This “*spill over*” effect of work stressors on mood at home was found to be greater than the “*spill over*” effect of stressors at home on mood at work [73]. This is consistent with the findings that when there are stressors both at work and at home, subjects believe their problems at home are in fact due to those at work [23]. Similarly, when assessed using a life events interview [75], both work and nonwork stressors contributed to depression in medical personnel. Even when vulnerability factors and nonwork stressors were controlled, work stress still contributed to depression.

The *pathways* linking work stressors and depression may not be direct since in a study of blue-collar workers [76] neither initial life events nor “work strain” (economic difficulties, work overload) directly predicted later depression: they did, however, predict subsequent life events and work strain, both of which correlated with depression at that time. This study also revealed a reciprocal relationship; initial depression also predicted subsequent life events and work strain [76].

#### Findings from intervention studies

Intervention studies can provide the most robust evidence of a causal relationship but they are few in this area. Briner [77] and Cooper and Cartwright [78] have reviewed stress interventions in the workplace and emphasise the need to distinguish *primary interventions* (organisation/structural change), *secondary interventions* (stress management/coping strategies) and *tertiary interventions* (interventions targeted for those actually stressed).

*Secondary and tertiary interventions* appear to have short-term effects only (in the order of 3 months), and indeed these interventions may be treating morbidity due to nonoccupational stressors [77,79]. Improvement may be seen in a range of psychological outcomes including psychological symptoms, self-esteem and perceptions of work stress [80].

Primary intervention studies report inconsistent findings. While job satisfaction may improve with ‘intervention’ [81–85], mental health may not [81,82,86], and absenteeism or staff turnover can indeed increase [81,82]. One recent study [87], however, found primary intervention was effective in reducing both depression and sick leave, despite a concurrent increase in amount of overtime worked. Another [79], comparing a primary intervention with (tertiary) individual stress management, found the former had no impact on mental health status (GHQ) and paradoxically was associated with increased physical symptoms, while the latter had only short-term positive effects consistent with other tertiary intervention studies [88]. There is thus very limited evidence that primary (organisational) intervention has significant mental health benefit while tertiary interventions may have positive but short-lived effects and may well be impacting on morbidity due to other nonoccupational causes.

#### Discussion

There are methodological limitations to many of these studies. Firstly, measures of psychological morbidity most commonly employed are self-report measures; rarely do the studies assess “psychiatric caseness” as diagnosed by a clinician, for obvious logistical reasons given the large sample sizes. Furthermore, other studies use “burnout” as an outcome measure which is not well validated as a construct, even though it does have some significant relationship to psychological health status, particularly depression [2,89]. Nonetheless, it may be a more relevant measure in assessing psychological distress in the workplace.

Secondly, work stressor variables are also often based on self-report. While this may seem to be a more “sensitive” method, it introduces potential bias due to “common method” of assessment, which can occur when a common perceptual set influences the reporting of both the “causal” and ‘outcome’ variables. For instance *subjective* appraisals

of work stress correlated more strongly with self-reported depression than did *objective work conditions* [16,90]; it is indeed argued that the relation between work stress and depression or burnout may be simply attributable to underlying career frustration [2]. While “self-report” methodology has limitations, they can be largely overcome with prospective study design and appropriate control of “time one” and other confounding covariates such as evidenced in the Whitehall II study [15–21,66–68].

Thirdly, some studies have shown that there may also be a “reverse causal” or reciprocal relationship; both burnout [33] and depression [46,76,91] can also effect the perception or experience of work stressors. Similarly, one study [92] has demonstrated a reciprocal relation between “job resources” (autonomy, variety, skill utilisation) and mental health. It is argued that multipanel prospective studies with “objective” work stressor indices, preferably using structural equation approaches, can best address this issue [93].

Fourthly, “third variables” (such as personality) are also crucial; their effects may be complex and are often poorly assessed. Personality firstly seems to be an independent risk factor for both depression and burnout, irrespective of exposure to work stressors [94]. Secondly, personality (if not appropriately controlled) may also confound the observed association between work stressors and depression since it can influence the perception or occurrence of stressors at work, and depression risk [90]. Finally, personality could moderate the stressor–depression relationship; however, the evidence is weak. When however these ‘third’ variables have been appropriately controlled in prospective studies, occupational stressors remain as independent predictors of depression.

Finally, in relation to broader issues of method, Mansella [95] argues for the application of a wide variety of methods, measures and sources of subjects in assessing the adverse effects of occupational stress while Kristensen [96] proposes that because stressors clearly differ from one occupation to another, that research therefore should be occupationally specific.

## Conclusions

Occupational stress is of increasing importance due to continuing structural changes in the workplace, with both increasing demands and job insecurity imposed on employees. A range of adverse health outcomes have been identified but psychological disorders are significant because they occur frequently, are often unrecognised and can be accompanied by significant social morbidity; there are important implications for the lifestyle and health of employees and their families and there are medico-legal issues.

The major medico-legal issue is that of “liability”; there are two important considerations in this regard. The first is to determine what degree of work stress (responsibilities, hours worked, etc.) may be likely to cause psychological impair-

ment in an average employee, but as yet no reliable quantification is available. In individual cases judgement still rests with experienced clinicians using their clinical findings, with research findings as an evidence-based guide. The problem of assessing what may be acceptable or unacceptable stress is further confused by virtue of the fact that increasing responsibilities and hours worked are becoming more common in the workplace and thus seem more “normative.”

The second consideration is, who takes responsibility for the stress exposure. Is it driven by the employer or employee? Employers rarely suggest their employees are working too hard and that they should “ease up”, indeed they generally expect increasing productivity from their work force because they are driven by shareholder profit expectations and because they believe perhaps their employees are sufficiently well-renumerated for their efforts. At the same time, the average employee accepts the situation because, although not explicitly stated, their continued employment may depend on meeting these high demands. Other employees, however, those who are workaholic and/or very ambitious, may themselves be more responsible for their exposure to work stress. While the relationship between the employer, employee and the determination of work stress may be complex, it seems clear that psychological morbidity in the workplace will continue as a significant problem unless major changes occur in the structure of employment.

## References

- [1] Hotopf M, Wessely S. Stress in the workplace: unfinished business. *J Psychosom Res* 1997;43(1):1–6.
- [2] Glass D, McKnight JD. Perceived control, depressive symptomatology, and professional burnout: a review of the evidence. *Psychol Health* 1996;11(1):23–48.
- [3] Maslach C. Burnout research in the social sciences: a critique. *J Soc Serv Res* 1987;10:95–105.
- [4] Iacovides A, Fountoulakis KN, Moysidou C, Lerodiakonou C. Burnout in nursing staff: is there a relationship between depression and burnout? *Int J Psychiatry Med* 1999;29(4):421–33.
- [5] Baba V, Galperin BL, Lituchy TR. Occupational mental health: a study of work-related depression among nurses in the Caribbean. *Int J Nurs Stud* 1999;36(2):163–9.
- [6] Karasek RA. Job demands, job decision latitude and mental strain: implications for job redesign. *Adm Q Sci* 1979;24:285–306.
- [7] Kawakami N, Haratani T, Araki S. Effects of perceived job stress on depressive symptoms in blue-collar workers of an electrical factory in Japan. *Scand J Work, Environ Health* 1992;18(3):195–200.
- [8] Kawakami N, Araki S, Kawashima M. Effects of job stress on occurrence of major depression in Japanese industry: a case-control study nested in a cohort study. *J Occup Med* 1990;32(8):722–5.
- [9] Aro S, Hasan J. Occupational class, psychosocial stress and morbidity. *Ann Clin Res* 1987;19(2):62–8.
- [10] Melamed S, Luz J, Green MS. Noise exposure, noise annoyance and their relation to psychological distress, accident and sickness absence among blue-collar workers — the Cordis Study. *Isr J Med Sci* 1992;28(8):629–35.
- [11] Stansfeld SA, Sharp DS, Gallacher J, Babisch W. Road traffic noise, noise sensitivity and psychological disorder. *Psychol Med* 1993;23(4):977–85.
- [12] Stansfeld SA. Noise, noise sensitivity and psychiatric disorder: epi-

- demological and psychophysiological studies. *Psychol Med* 1992; 22(Suppl. 22):1–44.
- [13] Romanov K, Appelberg K, Honkasalo ML, Koskenvuo M. Recent interpersonal conflict at work and psychiatric morbidity: a prospective study of 15,530 employees aged 24–64. *J Psychosom Res* 1996; 40(2):169–76.
  - [14] Borg V, Kristensen TS, Burr H. Work environment and changes in self-rated health: a five year follow-up study. *Stress Med* 2000;16(1): 37–47.
  - [15] Stansfeld SA, Fuhrer R, Head J, Ferrier J, Shipley M. Work and psychiatric disorder in the Whitehall II study. *J Psychosom Res* 1997;43(1):73–81.
  - [16] Stanfield SA, North FM, White I, Marmot MG. Work characteristics and psychiatric disorder in civil servants in London. *J Epidemiol Community Health* 1995;49(1):48–53.
  - [17] Stansfeld SA, Bosma H, Hemingway H, Marmot MG. Psychosocial work characteristics and social support as predictors of SF-36 health functioning: the Whitehall II study. *Psychosom Med* 1998;60(3): 247–57.
  - [18] Stansfeld SA, Head J, Marmot MG. Explaining social class differences in depression and well-being. *Soc Psychiatry Psychiatr Epidemiol* 1998;33(1):1–9.
  - [19] Ferrie JE, Shipley MJ, Marmot MG, Stansfeld SA, Smith GD. An uncertain future: the health effects of threats to employment security in white-collar men and women. *Am J Public Health* 1998;88(7): 1030–6.
  - [20] Ferrie JE, Shipley MJ, Marmot MG, Stansfeld S, Davey Smith G. The health effects of major organisational change and job insecurity. *Soc Sci Med* 1998;46(2):243–54.
  - [21] Ferrie JE, Shipley MJ, Marmot MG, Stansfeld SA, Smith GD. Health effects of anticipation of job change and non employment: longitudinal data from the Whitehall II study. *BMJ* 1995;311(7015):1264–9.
  - [22] Heinisch DA, Jex SM. Negative affectivity and gender as moderators of the relationship between work-related stressors and depressed mood at work. *Work Stress* 1997;11(1):46–57.
  - [23] Cherry N. Stress, anxiety and work: a longitudinal study. *J Occup Psychol* 1978;5(3):259–70.
  - [24] Frone MR, Russell M, Cooper ML. Job stressors, job involvement and employee health: a test of identity theory. *J Occup Organ Psychol* 1995;68(1):1–11.
  - [25] Parkes KR. Coping, negative affectivity, and the work environment: additive and interactive predictors of mental health. *J Appl Psychol* 1990;75(4):399–409.
  - [26] Schonfeld IS. Relation of negative affectivity to self-reports of job stressors and psychological outcomes. *J Occup Health Psychol* 1996;1(4):397–412.
  - [27] Schonfeld IS. A longitudinal study of occupational stressors and depressive symptoms in first-year female teachers. *Teach Teach Educ* 1992;8:151–8.
  - [28] Burke RJ, Greenglass E. A longitudinal study of psychological burnout in teachers. *Hum Relat* 1995;48(2):187–202.
  - [29] Burke RJ, Greenglass ER. A longitudinal examination of the Cherniss model of psychological burnout. *Soc Sci Med* 1995;40(10):1357–63.
  - [30] Burke RJ, Greenglass ER, Schwarzer R. Predicting teacher burnout over time: effects of work stress, social, and self-doubts on burnout and its consequences. *Anxiety, Stress Coping: Int J* 1996;9(3): 261–75.
  - [31] Brenner SO, Soerboom D, Wallius E. The stress chain: a longitudinal confirmatory study of teacher stress, coping and social support. *J Occup Psychol* 1985;58(1):1–13.
  - [32] Jackson S, Schwab R, Schuler R. Toward an understanding of the burnout phenomenon. *J Appl Psychol* 1986;61:630–40.
  - [33] Mills LB, Huebner ES. A prospective study of personality characteristics, occupational stressors, and burnout among school psychology practitioners. *J Sch Psychol* 1998;36(1):103–20.
  - [34] Kapur N, Borrill C, Stride C. Psychological morbidity and job satisfaction in hospital consultants and junior house officers: multicentre, cross sectional survey. *BMJ* 1998;317:511–22.
  - [35] Revicki DA, Whitley TW, Gallery ME, Allison EJ. The impact of work environment characteristics on work-related stress and depression in emergency medical residents: a longitudinal study. *J Community Appl Soc Psychol* 1993;3(4):273–84.
  - [36] Rout U, Cooper CL, Rout JK. Job stress among British general practitioners: predictors of job dissatisfaction and mental ill-health. *Stress Med* 1996;12(3):155–66.
  - [37] Cooper GL, Clarke S, Rowbottom AM. Occupational stress, job satisfaction and well-being in anaesthetists. *Stress Med* 2000;15(2): 115–26.
  - [38] Johnson JV, Hall EM, Ford DE, Mead LA, Levine DM, Wang NY, Klag MJ. The psychosocial work environment of physicians. The impact of demands and resources on job dissatisfaction and psychiatric distress in a longitudinal study of Johns Hopkins Medical School graduates. *J Occup Environ Med* 1995;37(9):1151–9.
  - [39] Wall TD, Bolden RI, Borrell CS, Carter AJ, Golya DA, Hardy GE, Haynes CE, Rick JE, Shapiro DA, West MA. Minor psychiatric disorder in NHS trust staff: occupational and gender differences. *Br J Psychiatry* 1997;171:519–23.
  - [40] Firth H, Britton P. “Burnout” absence and turnover amongst British nursing staff. *J Occup Psychol* 1989;62(1):55–9.
  - [41] Burke RJ, Greenglass ER. Work status congruence work outcomes and psychological well-being. *Stress Med* 2000;16(2):91–9.
  - [42] Wade DC, Cooley E, Savicki V. A longitudinal study of burnout. *Child Youth Serv Rev* 1986;8(20):161–73.
  - [43] Schaefer JA, Moos RH. Effects of work stressors and work climate on long-term care staff’s job morale and functioning. *Res Nurs Health* 1996;19(1):63–73.
  - [44] Poulin J, Walter C. Social work burnout: a longitudinal study. *Soc Work Res Abstr* 1993;29(4):5–11.
  - [45] Savicki V, Cooley EJ. Burnout in child protective service workers: a longitudinal study. *J Organ Behav* 1994;15(7):655–66.
  - [46] Prosser D, Johnson S, Kuipers E. Perceived sources of work stress and satisfaction among hospital and community mental health staff, and their relation to mental health, burnout and job satisfaction. *J Psychosom Res* 1997;43(1):51–9.
  - [47] Richman JA, Rospenda KM, Kawyn SJ, Flaherty JA, Fendrich M, Drum ML, Johnson TP. Sexual harassment and generalised workplace abuse among university employees: prevalence and mental health correlates. *Am J Public Health* 1999;89(3):358–63.
  - [48] Quine L. Workplace bullying in NHS community trust: staff questionnaire survey. *BMJ* 1999;318(7178):228–32.
  - [49] Lam LT, Ross FI, Cass DT, Quine S, Lazarus R. The impact of work related trauma on the psychological health of nursing staff: a cross sectional study. *Aust J Adv Nurs* 1999;16(3):14–20.
  - [50] Lowenthal KM, Eysenck M, Harris D, Lubitsh G, Gorton T, Bicknell H. Stress, distress and air traffic incidents: job dysfunction and distress in airline pilots in relation to contextually-assessed stress. *Stress Med* 2000;16(3):179–83.
  - [51] Kessler RC. The effects of stressful life events on depression. *Annu Rev Psychol* 1997;48:191–214.
  - [52] Spurgeon A, Harrington MJ, Cooper CL. Health and safety problems associated with long working hours: a review of the current position. *Occup Environ Med* 1997;54:367–75.
  - [53] Bhese PD, Halverson RR. Individual, nomothetic models of job stress. An examination of work hours, cohesion and well-being. *J Appl Soc Psychol* 1996;26:1171–81.
  - [54] Watanabe S, Tori J, Shinkai S, Watanabe T. Relationships between health status and working conditions and personalities among VDT workers. *Environ Res* 1993;61:258–65.
  - [55] Hall E, Johnson JV, Stewart W. Double exposure: the combined impact of the home and work environs on psychosomatic strain in Swedish women and men. *Int J Health Serv* 1992;22:239–45.
  - [56] Turnage JJ, Spielberger CD. Job stress in managers, professional and clerical workers. *Work Stress* 1991;5:165–76.
  - [57] Burke RJ. Workaholicism in organizations: psychological and physical well-being consequences. *Stress Med* 2000;16(1):11–6.

- [58] Batlis NC. Job involvement and locus of control as moderators of role-perception/individual-outcome relationships. *Psychol Rep* 1980; 46:111–9.
- [59] Watson DC. Negative affectivity: the disposition to experience aversive emotional states. *Psychol Bull* 1984;96(3):465–90.
- [60] Brett MJ, Brief AP, George JM. The role of negative affectivity in understanding relations between self-reports of stressors and strains: a comment on the applied psychology literature. *J Appl Psychol* 1993; 78(3):402–12.
- [61] Burke JF, Brief AP. Negative affectivity and the reporting of stressful life events. *Health Psychol* 1990;9(1):57–68.
- [62] Spector PE, O'Connell BJ. The contribution of personality traits, negative affectivity, locus of control and Type A to the subsequent reports of job stressors and job strains. *J Occup Organ Psychol* 1994; 67(1):1–12.
- [63] Necowitz LB, Roznowski M. Negative affectivity and job satisfaction: cognitive processes underlying the relationship and effect on employee behaviours. *J Vocational Behav* 1994;45(3):270–94.
- [64] Krantz G, Ostergren PO. Common symptoms in middle aged women: their relation to employment status, psychosocial work conditions and social support in a Swedish setting. *J Epidemiol Community Health* 2000;54(3):192–9.
- [65] Nadaoka T, Kashiwakura M, Oiji A, Morioka Y, Totsuka S. Stress and psychiatric disorders in local government officials in Japan, in relation to their employment level. *Acta Psychiatr Scand* 1997;96(3):176–83.
- [66] Stansfeld SA, Fuhrer R, Shipley MJ. Types of social support as predictors of psychiatric morbidity in a cohort of British civil servants (Whitehall II study). *Psychol Med* 1998;28(4):881–92.
- [67] Stansfeld SA, Rael EG, Head J, Shipley M, Marmot M. Social support and psychiatric sickness absence: a prospective study of British civil servants. *Psychol Med* 1997;27(10):35–48.
- [68] Rael EGS, Stansfeld SA, Shipley M, Head J, Feeney A, Marmot M. Sickness absence in the Whitehall II study. London: the role of social support and material problems. *J Epidemiol Community Health* 1995;49(5):474–81.
- [69] Frese M. Social support as a moderator of the relationship between work stressors and psychological dysfunctioning: a longitudinal study with objective measures. *J Occup Health Psychol* 1999;4(3):179–92.
- [70] Hagihara A, Tarumi K, Miller AS. Social support at work as a buffer of work stress-strain relationship: a signal detection approach. *Stress Med* 1998;14(2):75–81.
- [71] Dormann C, Zapf D. Social support, social stressors at work, and depressive symptoms: testing for main and moderating effects with structural equations in a three-wave longitudinal study. *J Appl Psychol* 1999;84(6):874–84.
- [72] Phelan J, Schwartz JE, Bromet EJ, Dew MA, Parkinson DK, Schulberg HC, Dunn LO, Blane H, Curtis EC. Work stress, family stress and depression in professional and managerial employees. *Psychosom Med* 1991;21(4):999–1012.
- [73] Leiter MP, Durup MJ. Work, home, and in-between: a longitudinal study of spillover. *J Appl Behav Sci* 1996;32(1):29–47.
- [74] Noor NM. Work and family roles in relation to women's well-being: a longitudinal study. *Br J Soc Psychol* 1995;34(Pt 1):87–106.
- [75] Weinberg A, Creed F. Stress and psychiatric disorder in health care professionals and hospital staff. *Lancet* 2000;355(9203):533–7.
- [76] Glickman L, Tanaka JS, Chan E. Life events, chronic strain and psychological distress: longitudinal causal models. *J Community Psychol* 1991;19(4):283–305.
- [77] Briner RB. Improving stress assessment: toward an evidence-based approach to organizational stress interventions. *J Psychosom Res* 1997;43(1):61–71.
- [78] Cooper CL, Cartwright S. An intervention strategy for workplace stress. *J Psychosom Res* 1997;43(1):7–16.
- [79] Reynolds S. Psychological well-being at work: is prevention better than cure? *J Psychosom Res* 1997;43(1):93–102.
- [80] Firth-Gozens JA, Hardy GE. Occupational stress, clinical treatment and changes in job perceptions. *J Occup Organ Psychol* 1992;65: 81–8.
- [81] Wall TD, Kemp NJ, Jackson PR, Paul R, Clegg CW. Outcomes of autonomous workgroups: a long-term field experiment. *Acad Manage J* 1986;29: 280–304.
- [82] Gordery JL, Mueller WS, Smith LM. Attitudinal and behavioural effects of autonomous group working: a longitudinal study. *Acad Manage J* 1991;34:464–76.
- [83] Cooper CI, Williams S. Creating healthy work organizations. Chichester: Wiley, 1997.
- [84] Quick JC. Dyadic goal setting and role stress in a field study. *Acad Manage J* 1979;22:241–52.
- [85] Jackson SE. Participation in decision making as a strategy for reducing job related strain. *J Appl Psychol* 1983;68:3–19.
- [86] Heaney CA, Isreal BA, Schurman SJ, Baker EA, House JS, Hugentobler M. Industrial relations, worksite stress reduction and employee well-being: a participatory action reinvestigation. *J Organ Behav* 1993;14:495–510.
- [87] Kawakami N, Araki S, Kawashima M, Masumoto T, Hayashi T. Effects of work-related stress reduction on depressive symptoms among Japanese blue-collar workers. *Scand J Work, Environ Health* 1997;23(1):54–9.
- [88] Murphy LR. Stress management in highway maintenance workers. *J Occup Med* 1984;26(6):436–42.
- [89] Golembiewski RT, Munzenrider RF, Scherb K, Billingsley W. Burn-out and psychiatric "cases": early evidence of an association. *Anxiety, Stress Coping: Int J* 1992;5(1):69–78.
- [90] Goldberg P, David S, Landre MF, Goldberg M, Dassa S, Fuhrer R. Work conditions and mental health among prison staff in France. *Scand J Work, Environ Health* 1996;22(1):45–54.
- [91] Daniels K, Guppy A. Stressors, locus of control, and social support as consequences of affective psychological well-being. *J Occup Health Psychol* 1997;2(2):156–74.
- [92] Taris TW. The mutual effects between job resources and mental health: a prospective study among Dutch youth. *Genet, Soc Gen Psychol Monogr* 1999;125(4):433–50.
- [93] Zapf D, Dormann C, Frese M. Longitudinal studies in organizational stress research: a review of the literature with reference to methodological issues. *J Occup Health Psychol* 1996;1(2):145–69.
- [94] Taylor H, Cooper CL. The stress-prone personality: a review of the research in the context of occupational stress. *Stress Med* 1989;5: 17–27.
- [95] Mansella AJ. The measurement of emotional reactions to work: conceptual methodological and research issues. *Work Stress* 1994;8(2): 153–76.
- [96] Kristensen TS. Job stress and cardiovascular disease: a theoretic critical review. *J Occup Health Psychol* 1996;1(3):246–60.